



- I.** The first coordinate system shows the graph of a function $f(x)$. On the second coordinate system, sketch the graph of its derivative $f'(x)$.
(4)



- II.** The portion of the graph of a function $g(x)$ for $0 \leq x \leq 1$ is sketched on the first coordinate system below. The second coordinate system shows the graph of the derivative $g'(x)$. On the first coordinate system, sketch the remaining part of the graph of $g(x)$.
(4)



- III.** Use the definition of limit to prove that $\lim_{x \rightarrow 3} 7x + 1 = 22$.
(4)

x	$f(x)$	$g(x)$	$f'(x)$	$g'(x)$
1	-1	2	2	3
2	1	8	3	0
3	5	2	1	9

IV. A table giving some of the values for f , g , f' , and g' is given here:
(12)

For each of the following questions, please **circle the letter** of the correct answer, based on the values of f and g given in the table.

- If $h(x) = f(g(x))$, find $h'(1)$.
 - 3
 - 0
 - 2
 - 3
 - 5
 - 6
 - 9
 - 12
 - 15
- If $f''(1) = 1$, $f''(2) = 4$, and $f''(3) = 2$, what is the value of the derivative of $f(f'(x))$ at $x = 2$?
 - 2
 - 3
 - 4
 - 6
 - 8
 - 12
 - 16
 - 20
 - 21
- Based on the values given in the table, the linear part of the change of g as x changes from 2 to 3 is:
 - 6
 - 1
 - 0
 - 3
 - 4
 - 5
 - 6
 - 8
 - 9
- If f is differentiable, then (based on the values of f given in the table) the Mean Value Theorem guarantees that $f'(x)$ must assume which one of the following values:
 - 7
 - 1
 - 6
 - 1
 - 12
 - 4
 - 0
 - 18
 - 21
- If f is continuous, then it must have a root between $x = 0$ and $x = 1$. One can conclude this using which of the following?
 - The Second Derivative Test
 - The Extreme Value Theorem
 - The First Derivative Test
 - Newton's Method
 - Rolle's Theorem
 - The Mean Value Theorem
 - The Fundamental Theorem of Algebra
 - The Intermediate Value Theorem
 - Linear Approximation
- If f is continuous, then it must assume a minimum value on the closed interval $[0, 1]$. One can conclude this using which of the following?
 - The Second Derivative Test
 - The Extreme Value Theorem
 - The First Derivative Test
 - Newton's Method
 - Rolle's Theorem
 - The Mean Value Theorem
 - The Fundamental Theorem of Algebra
 - The Intermediate Value Theorem
 - Linear Approximation

V. For each of the following, find *all* $f(x)$ that satisfy the given condition or conditions.

(9)

1. $f'(x) = 3x^2 - \pi x^3$

2. $f'(x) = 3 + \sin(x)$ and $f(0) = 2$

3. $f''(x) = x$

VI. Give a precise mathematical definition of each of the following:

(7)

1. $f(x)$ has a *local minimum* at $x = a$ (use an open interval to express the idea that x is near a)

2. $x = a$ is a *critical point* (also called a *critical number*) of $f(x)$

3. $\lim_{f \rightarrow L} k(f) = a$ (give the ϵ - δ definition)

VII. For each of the following questions, please **circle the letter** of the correct answer,
(8)

1. Suppose that L is a function for which $L'(x) = \frac{1}{x}$. The derivative of $(L(x))^2$ is:

- | | | |
|--------------------------|-------------------------|-----------------------|
| (a) $\frac{2}{L(x)}$ | (b) $2xL(x)$ | (c) $\frac{2L(x)}{x}$ |
| (d) $-\frac{2L(x)}{x^2}$ | (e) $\frac{2L(x)}{x^2}$ | (f) $\frac{2}{x}$ |
| (g) $2L(x)$ | (h) $\frac{1}{x^2}$ | (i) $-\frac{1}{x^2}$ |


2. Which of the following can be applied to a function *only* when the function is a polynomial?

- | | | |
|--|------------------------------------|-------------------------------|
| (a) The Second Derivative Test | (b) The Extreme Value Theorem | (c) The First Derivative Test |
| (d) Newton's Method | (e) Rolle's Theorem | (f) The Mean Value Theorem |
| (g) The Fundamental Theorem of Algebra | (h) The Intermediate Value Theorem | (i) Linear Approximation |

3. The derivative of a certain function r is $r'(x) = \frac{x}{1+x^2}$. At $x = 0$, r has

- | | | |
|-------------------------|--|--|
| (a) an absolute minimum | (b) a local minimum which is not an absolute minimum | (c) a point where the graph of r has vertical slope |
| (d) an absolute maximum | (e) a local maximum which is not an absolute maximum | (f) a critical point where r has neither a local maximum nor a local minimum |
| (g) an inflection point | (h) a vertical asymptote | (i) a horizontal asymptote |

VIII. As shown in the figure to the right, a ladder 10 feet long leans against a wall. Its base is sliding directly away from the wall at 2 feet per second. How fast is the angle θ changing when θ is $\pi/4$?



IX. Calculate the following derivatives.


(12)

1. $\frac{dy}{dx}$ if $y^2 = \cos(x + y)$ (use implicit differentiation)

2. $\frac{(x^3 + 7)^4}{\sin(x)}$

3. $f''(x)$ if $x = \tan(x)$

- X.** The figure to the right shows an equilateral triangle with sides of length L and a rectangle inscribed in the triangle. Express the area of the rectangle in terms of the x -coordinate of the lower right-hand corner of the rectangle. Do *not* proceed further with the maximization problem, just give the area A as a function of x and no other variables.
- (4)



- XI.** The figure to the right shows a rectangular box with open top. The base has length twice its width. The volume of the box is 5 cubic units. We let x be the width of the base, so that the length of the base is $2x$ and the height is $\frac{5}{2x^2}$. Calculate the surface area S (the total of the 4 sides, plus the base) in terms of x , and find the value of x that minimizes S .
- (6)

